

# Sipcafol Tokio

## EC FERTILISER

NPK fertiliser blend 9-54-6 with micro-nutrients

### COMPOSITION

Total nitrogen (N).....	9%
Ammoniacal nitrogen (NH4).....	8,5%
Nitric Nitrogen.....	0,5%
Phosphorus pentaoxide (P <sub>2</sub> O <sub>5</sub> ) total soluble in water and ammonium citrate.....	54%
Potassium oxide (K <sub>2</sub> O) soluble in water (low chloride content).....	6%
Iron (Fe) soluble in water.....	0.1%
Total Iron (Fe) complexed by EDTA.....	0.1%
Zinc (Zn) soluble in water.....	0.05%
Total zinc (Zn) complexed by EDTA.....	0.05%
Manganese (Mn) soluble in water.....	0.05%
Total manganese (Mn) complexed by EDTA.....	0.05%
Copper (Cu) soluble in water .....	0.05%
Total Copper (Cu) complexed by EDTA.....	0.05%
Boron (B) soluble in water .....	0.5%
Iron chelated by EDTA is stable in the pH range 3-7. Copper, manganese, and zinc chelated by EDTA are stable in the pH range 4-11.	

**Net weight: 5 Kg**

**FORMULATION:** soluble microcrystals

**Producer:** SIPCAM Inagra - Calle Prof Beltran  
Baguena,5 46009 Valencia Spagna  
Tel. +34 (963) 483 500

### Danger/Risks/Caution

P102 Keep out of reach of children and of incompetent people. P270 Do not eat, drink or smoke during use. P264 Wash hands thoroughly after handling. P280 Gloves / clothing Protective / eye protection / face protection. P302 + P352 IF ON SKIN: Wash with plenty of soap and water. P305 + P351 + P338 IN CASE OF EYE CONTACT: Rinse thoroughly for several minutes. Remove contact lenses if easy to do. Continue rinsing. P233 Keep container tightly closed

### ADDITIONAL REQUIREMENTS

Do not contaminate water with the product or the container

### Dosage and usage indications

#### Foliar application

CROPS	TIMINGS	Nº appl	RATES (Lt/Ha)
Orchard	From Pre-flowering to Setting	2-3 every 15-20 days	2-4
Vegetables	From Pre-flowering to Setting	4-5 every 15-20 days	2-4
Flowers and Ornamentals	From pre-flowering	2-5 every 15-20 days	150-200 g/100 L water
Oil crops (Oilseed rape, Sunflower, Soybean, Peas)	From Pre-flowering to Setting	1-2	2-3
Potato	From Pre-flowering to Tuber Development	2-3 every 15-20 days	3-4

**WARNING: In case of low volumes, do not exceed 500 g/hl**

**Application and dosages vary according to soil characteristics, crop growth stage and pedoclimatic conditions. Please consult the Technical Service for application advice in specific crop and pedoclimatic conditions. The product is compatible with almost all the insecticides and fungicides to the exclusion of formulations containing copper, mineral oils and emulsion but given the wide variety of products on the market in the case of mixture perform a preliminary test of compatibility on small surfaces . SIPCAM responds only about the composition of the product. Instead assumes no responsibility for any damage caused by misuse and / or abuse of the product.**

### WARNINGS

**Keep away from open flames, in a cool and breezy warehouse. Avoid exposure to low temperatures for long periods. Periodically check the packaging. In case of accidental spillage stem losses, avoid contamination of waterways and, if necessary, dispose of according to applicable regulations. Shake well before use. Dispose of packaging according to applicable regulations. The container must not be emptied completely dispersed in the environment. The container can not be reused.**