

# Fezan<sup>®</sup> Plus

MAPP 15490

A suspension concentrate containing 60 g/l (5.45% w/w) tebuconazole and 166 g/l (15.1% w/w) chlorothalonil.

A fungicide for disease control in Winter Wheat.

## WARNING



- Harmful if inhaled
- Suspected of causing cancer



- Suspected of damaging the unborn child
- Very toxic to aquatic life with long lasting effects



- Avoid breathing spray.
- Wear suitable protective clothing (coveralls), suitable protective gloves and face protection (faceshield) when handling the concentrate and contaminated surfaces.
- If inhaled; remove victim to fresh air and keep at rest in a position comfortable for breathing.
- If exposed or concerned; get medical advice/attention.
- Collect spillage.
- Avoid release to the environment.
- To protect aquatic organisms respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.
- Dispose of contents/container to a licensed waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.
- Do not contaminate water with the product or its container (do not clean application equipment near surface water / avoid contamination via drains from farmyards and roads).

Contains chlorothalonil. May produce an allergic reaction. Contains 1,3,5-tris (2-hydroxyethyl) hexahydro-1,3,5-triazine. May cause an allergic reaction. To avoid risks to man and the environment, comply with the instructions for use. The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work. FOR 24 HOUR EMERGENCY INFORMATION TELEPHONE **01763 212 100**.

### Registration Holder

OXON ITALIA S.P.A.  
Via Sempione 151,  
20016 Pero (MI) Italy

### Marketing Company

SIPCAM UK Ltd, 4C Archway House,  
The Lanterns, Melbourn Street,  
Royston, Herts SG8 7BX

FEZAN<sup>®</sup> PLUS is a registered trademark and product of OXON ITALIA SpA



Scan to view the Safety Data Sheet

Alternatively, download the Safety Data Sheet from [sipcamuk.co.uk](http://sipcamuk.co.uk) or contact your supplier.

[sipcamuk.co.uk](http://sipcamuk.co.uk)

# FUNGICIDE

Net contents:

# 10Le

  
**SIPCAM**  
UK

## IMPORTANT INFORMATION

### FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crops	Maximum individual dose (L product/hectare)	Maximum total dose (L product / ha / crop)	Latest time of application
Winter Wheat	3.0	6.0	Before end of flowering (BBCH 69)

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

## SAFETY PRECAUTIONS

### Operator Protection

Engineering control of operator exposure must also be used where reasonably practicable in addition to the following personal protective equipment.

Operators must wear suitable protective gloves when handling the concentrate or handling contaminated surfaces.

Do not breathe spray.

When using do not eat, drink or smoke.

Wear suitable protective clothing and gloves.

However, other engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

IF YOU FEEL UNWELL seek medical advice immediately (show the label where possible).

### Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

DO NOT ALLOW DIRECT SPRAY from horizontal boom



sprayers to fall within 5m of the top of the bank of a static or flowing waterbody, or within 1m of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

### Storage and Disposal

- Keep out of reach of children.
- Keep away from food, drink and animal feedingstuffs.
- Use appropriate containment to avoid environmental contamination.
- This material and its container must be disposed in a safe way.
- KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.
- RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.
- DO NOT RE-USE CONTAINER for any purpose.
- Protect from frost.

## DIRECTIONS FOR USE

---

### IMPORTANT

This information is authorised as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

### Resistance

FEZAN® PLUS contains a DMI fungicide. Resistance to some DMI fungicides has been identified in Septoria leaf blotch (*Mycosphaerella graminicola*), which may seriously affect the performance of some products. For further advice on resistance management in DMI's, contact your agronomist or specialist advisor, and visit the Fungicide Resistance Action Group (FRAG)-UK website.

As part of a resistance management strategy non-chemical control measures, including the selection of resistant varieties, should be considered. Multiple applications of DMI fungicides made to the same crop to control diseases such as Septoria leaf blotch and Powdery mildew should be avoided. Where additional fungicide treatments are required to control disease throughout the season, products with a different (non-DMI) mode of action should be selected.

Consult processor before using on crops grown for processing

### Diseases Controlled

FEZAN® PLUS is intended for the use as a spray on Wheat (Winter) for protection and control of a range of wheat diseases including Septoria Leaf Blotch, Fusarium Ear Blight and Sooty Moulds.

FEZAN® PLUS is a mixture of fungicide active ingredients with different characteristics, combining the systemic protective, action of tebuconazole with the protectant activity of chlorothalonil.

### Crop Specific Information

All varieties of Winter Wheat may be treated with FEZAN® PLUS up to the end of flowering (BBCH 69).

DOSE: Apply FEZAN® PLUS at 3.0 L/hectare in 200-400 litres

of water per hectare. Apply as a MEDIUM quality spray as defined in the BCPC system.

Maximum total dose per crop; 6.0 l/ha.

A minimum of 7 days must be observed between applications.

TIMING: Moderate control of Septoria Leaf Blotch;

When Septoria Leaf Blotch is present at high levels early in the spring, and where the risk of disease development is high (e.g on susceptible varieties), an early treatment (e.g at the start of stem elongation) may be beneficial in reducing the spread and development of the disease.

A programme of two treatments applied during stem elongation (e.g. when leaf 3 is emerged) and at flag leaf emerged (e.g. flag leaf fully unrolled) will provide the best protection of the upper leaves and the ear. Apply when disease is active but before it appears on the upper leaves. Applications made when foliar symptoms are already present on the upper leaves will be less effective.

Control of Fusarium Ear Blight and Sooty Moulds;

Apply FEZAN® PLUS shortly after ear emergence (BBCH 59-69).

### Following Crops

There are no restrictions on following crops.

### Mixing and Spraying

Shake well before use. Half fill the spray tank with clean water and add the required quantity of FEZAN® PLUS while maintaining constant agitation. Add the remaining water and continue agitation during spray application.

## SIPCAM UK LTD CONDITIONS OF SUPPLY

---

All goods supplied by us are of high grade, but as we cannot exercise control over their mixing or use, all conditions or warranty, statutory or otherwise, as to the quality or fitness of our goods for any purpose are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.