

# Rover 500

Suspension concentrate containing 500 g/l (40.58% w/w) chlorothalonil.

A multi-site protectant fungicide for the control of fungal diseases on Wheat, Barley, Rye, Triticale and Field Beans.

## RISK AND SAFETY INFORMATION

### DANGER



- Harmful if inhaled
- Causes skin irritation
- Causes serious eye damage
- May cause an allergic skin reaction

- Suspected of causing cancer
- May cause respiratory irritation
- Very toxic to aquatic life with long lasting effects

- Avoid breathing spray.
- Do not get in eyes, on skin or on clothing.
- Wash hands and face thoroughly after handling.
- Wear gloves, protective clothing and eye/face protection.
- IF IN EYES, Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- Avoid release to the environment.
- Collect spillage.

- Dispose of contents/container to a licensed waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.
- Do not contaminate water with the product or its container (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads).
- To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

Contains chlorothalonil. May produce an allergic reaction.

**To avoid risks to human health and the environment, comply with the instructions for use.**

FOR 24 HOUR EMERGENCY INFORMATION TELEPHONE **+44(0)1763 212100**.

PCS Number 04467

#### Registration Holder

OXON ITALIA S.P.A.  
Via Sempione 195,  
20016 Pero (MI) Italy

#### Marketing Company

SIPCAM UK Ltd, 4C Archway House,  
The Lanterns, Melbourn Street,  
Royston, Herts SG8 7BX

**ROVER 500** is a registered trademark and product of OXON ITALIA SpA



Scan to view the Safety Data Sheet

Alternatively, download the Safety Data Sheet from [sipcamuk.co.uk](http://sipcamuk.co.uk) or contact your supplier.

[sipcamuk.co.uk](http://sipcamuk.co.uk)

# FUNGICIDE

Net contents:

# 10Le

  
**SIPCAM**  
UK

## FOR PROFESSIONAL USE ONLY

### DIRECTIONS FOR USE FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crops	Maximum individual dose (L product/hectare)	Maximum total dose (L product / ha / crop)	Latest time of application
Winter Wheat, Spring Wheat, Durum Wheat, Rye & Triticale	2.0	4.0	Before grain watery ripe stage (GS71)
Winter Barley & Spring Barley	2.0	4.0	Before beginning of flowering/first anthers visible (GS61)
Field Beans	2.0	4.0	Between, flowers open on first raceme (GS61) and, end of flowering (GS69)

**Other Specific Restrictions:** The product must only be used on beans that will be used as an animal feed item.

READ ALL PRECAUTIONS BEFORE USE

## SAFETY PRECAUTIONS

### Operator Protection

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD).

WASH SPLASHES FROM SKIN OR EYES IMMEDIATELY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

AVOID CONTACT BY MOUTH, WITH SKIN AND EYES.

WASH HANDS AND EXPOSED SKIN BEFORE EATING, DRINKING OR SMOKING AND AFTER WORK.

### Storage and Disposal

- KEEP OUT OF REACH OF CHILDREN.
- KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.
- KEEP IN ORIGINAL CONTAINER, TIGHTLY CLOSED, IN A SAFE PLACE.
- WASH OUT CONTAINER THOROUGHLY, EMPTY WASHINGS INTO SPRAY TANK AND DISPOSE OF SAFELY.
- THIS MATERIAL AND ITS CONTAINER MUST BE DISPOSED OF IN A SAFE WAY.
- USE APPROPRIATE CONTAINMENT TO AVOID ENVIRONMENTAL CONTAMINATION.

## DIRECTIONS FOR USE

---

### Mixing

To ensure even distribution of contents immediately before measuring, gently invert the container a few times. Half fill the spray tank with clean water and add the measured quantity of ROVER 500 whilst agitation is in progress. Complete filling of the water and continue agitation throughout the spraying application.

### Spray Quality

Apply as a **medium spray** as defined by **BCPC**.

### Cleaning

On emptying the container, rinse container thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

Thoroughly rinse the sprayer and measuring equipment immediately after use.

## SIPCAM UK LTD CONDITIONS OF SUPPLY

---

All goods supplied by us are of high grade, but as we cannot exercise control over their mixing or use, all conditions or warranty, statutory or otherwise, as to the quality or fitness of our goods for any purpose are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

## APPLICATION RATES AND TIMING

---

Please see next page.

## RESISTANCE MANAGEMENT

---

Consult your adviser for up to date guidance regarding current resistance status and a strategy for preventing and managing resistance in the cereal pathogens listed on the label. The Fungicide Resistance Action Committee (FRAC) produces recommendations that may be consulted for additional information.

## APPLICATION RATES AND TIMING

Crops	Diseases controlled	Timing	Application rate	Application volume
Winter and Spring Wheat, Durum Wheat, Rye and Triticale	<p>Modrate control:</p> <p>Leaf spot (<i>Septoria tritici</i>) and Glume blotch (<i>Leptosphaeria</i> (syn. <i>Septoria</i>) <i>nodorum</i>).</p> <p><b>Note:</b> Activity against Septoria may be reduced in the presence of serious mildew infection.</p>	<p>ROVER 500 contains chlorothalonil, a protectant fungicide best applied before disease onset or at the first sign of an attack.</p> <p>A single application in the early spring at growth stage 30-32 may prevent the spread of the disease within the crop.</p> <p>A later application from growth stage 33-70 may be used to protect the flag leaf and ear. For best results apply between growth stages 33-39.</p>	Apply ROVER 500 at 2 litres per hectare	200 to 400 litres of water per hectare as a MEDIUM spray (as defined by BCPC) at a pressure of at least 2.0 bar
Winter Barley, Spring Barley, Rye and Triticale	<p>Modrate control:</p> <p>Leaf scald (<i>Rhynchosporium secalis</i>)</p>	<p>ROVER contains chlorothalonil, a protectant fungicide best applied before disease onset or at the first sign of an attack.</p> <p>A single application as soon as disease is first detected in the crop.</p> <p>A second application may be made before growth stage 61 (before beginning of flowering/first anthers visible) if severe <i>Rhynchosporium</i> pressure occurs.</p>	Apply ROVER 500 at 2 litres per hectare	200 to 400 litres of water per hectare as a MEDIUM spray (as defined by BCPC) at a pressure of at least 2.0 bar
Field Beans	<p>Control of:</p> <p>Chocolate Spot (<i>Botrytis</i>)</p>	<p>Apply at early to mid flower and repeat two to three weeks later. Always make the first application before or as soon as disease is first seen and before it enters the 'aggressive' phase of infection. A two spray programme is essential.</p> <p>The product must only be applied to field beans between BBCH61 (flowers open on first raceme) and BBCH69 (end of flowering). The product must only be used on beans that will be used as an animal feed item.</p>	Apply ROVER 500 at 2 litres per hectare	200 to 400 litres of water per hectare as a MEDIUM spray (as defined by BCPC) at a pressure of at least 2.0 bar

Shake well before use. Protect from frost.