

Raikiri

(MAPP 20553)

A suspension concentrate containing 100 g/litre mesotrione.
A selective herbicide for the control of annual broad-leaved weeds
and barnyard grass in grain and forage maize.

RISK & SAFETY INFORMATION

24 HOUR EMERGENCY NUMBER: +44(0)1763 212100

DANGER



- May cause an allergic skin reaction
 - Causes serious eye damage
 - Very toxic to aquatic life with long lasting effects
 - Suspected of damaging the unborn child
-
- Obtain special instructions before use.
 - Avoid breathing spray.
 - Wear protective gloves/protective clothing/eye protection/face protection.
 - IF ON SKIN: Wash with plenty of soap and water.
 - IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE or doctor/physician.
 - If skin irritation or rash occurs: Get medical advice/attention.
 - Collect spillage.
 - Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Raikiri – contains 100 g/litre (9.4% w/w) mesotrione.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT: The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work

Authorisation Holder

Sumi Agro Europe Ltd
68 Upper Thames St, London.
EC4V 3BJ

Marketing Company

SIPCAM UK Ltd, 4C Archway House,
The Lanterns, Melbourn Street,
Royston, Herts SG8 7BX

sipcamuk.com

Tel +44(0)1763 212100
Fax +44(0)1763 212101

Net contents:

5Le

Batch number: SEE PACKAGE



Scan to view the Safety Data Sheet

Alternatively, download the Safety Data Sheet
from sipcamuk.co.uk or contact your supplier.

PROTECT FROM FROST.
SHAKE WELL BEFORE USE.

UFI: EESN-9CM4-3A0V-YQA2

sipcamuk.co.uk

SIPCAM
UK

HERBICIDE

MIXING

Half fill the sprayer tank with clean water and begin gentle agitation. Add the required quantity of product into the sprayer tank and allow to disperse fully. Rinse the empty containers thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add the washings to the sprayer and continue agitation whilst topping up the tank with water to the required level. Continue agitation until the mix is sprayed out. Spray immediately after mixing, do not allow the mixture to stand in the sprayer. When using tank-mixes follow any specified order of addition on the partner product label(s).

FOLLOWING CROPS

Crops following normal harvest of the treated crop:

AUTUMN-SOWN CROPS

Rye-grass, winter wheat, durum wheat, and winter barley can be planted in the Autumn following harvesting of a treated crop. Oilseed rape can be planted in the Autumn following harvesting of a treated crop, provided that the land is ploughed to a depth of at least 15cm and then cultivated prior to sowing.

SPRING-SOWN CROPS

Spring barley, spring wheat, rye-grass, forage maize and grain maize can be planted in the Spring following harvesting of a treated crop. DO NOT sow any other crop in the year following harvesting of a treated crop.

SIPCAM UK CONDITIONS OF SUPPLY

All goods supplied by us are of high grade, but as we cannot exercise control over their mixing or use, all conditions or warranty, statutory or other-wise, as to the quality or fitness of our goods for any purpose are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

FAILED CROPS

In the event of failure of a treated crop for any reason, plough prior to re-seeding. Sow only forage maize or grain maize after the failure of a treated crop. Some effect on the re-seeded crop may be seen shortly after emergence but these are usually only temporary and quickly outgrown.

COMPATIBILITY

Raikiri may be applied as a tank-mix with a range of products. Contact Sipcarn UK Ltd on 01763 212100 for compatibility information on specific tank-mixes.

SPRAYER DECONTAMINATION

Immediately after use thoroughly clean the sprayer and all equipment as follows: fill the spray tank with clean water to a minimum of 10% of its total volume. Flush the water through the sprayer, including nozzles and spray booms and lines. Drain and repeat the procedure. It is essential that all nozzles, filters, tubing, strainers, pumps and the spray tank are thoroughly cleaned after use to avoid the risk of damage to crops subsequently treated with the same equipment.



A suspension concentrate containing 100 g/litre mesotrione.
A selective herbicide for the control of annual broad-leaved weeds and barnyard grass in grain and forage maize.

RISK & SAFETY INFORMATION

24 HOUR EMERGENCY NUMBER: +44(0)1763 212100

DANGER



- May cause an allergic skin reaction
 - Causes serious eye damage
 - Very toxic to aquatic life with long lasting effects
 - Suspected of damaging the unborn child
-
- Obtain special instructions before use.
 - Avoid breathing spray.
 - Wear protective gloves/protective clothing/eye protection/face protection.
 - IF ON SKIN: Wash with plenty of soap and water.
 - IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE or doctor/physician.
 - If skin irritation or rash occurs: Get medical advice/attention.
 - Collect spillage.
 - Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Raikiri – contains 100 g/litre (9.4% w/w) mesotrione.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT: The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work

Authorisation Holder

Sumi Agro Europe Ltd
68 Upper Thames St, London.
EC4V 3BJ

Marketing Company

SIPCARN UK Ltd, 4C Archway House,
The Lanterns, Melbourn Street,
Royston, Herts SG8 7BX

sipcarnuk.com

Tel +44(0)1763 212100
Fax +44(0)1763 212101



Scan to view the Safety Data Sheet

Alternatively, download the Safety Data Sheet from sipcarnuk.co.uk or contact your supplier.

PROTECT FROM FROST.
SHAKE WELL BEFORE USE.

UFI: EESN-9CM4-3A0V-YQA2

sipcarnuk.co.uk

GROUP 27 HERBICIDE

HERBICIDE

Net contents:

5Le

Batch number: SEE PACKAGE



IMPORTANT INFORMATION FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Crops/situations:	Maximum individual dose	Maximum number of treatments	Latest time of application
Grain maize or forage maize	1.5 L/ha	1	8 leaves unfolded stage (GS18)

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

OTHER RESTRICTION: DO NOT HANDLE TREATED CROPS FOR AT LEAST 2 DAYS AFTER TREATMENT

GENERAL INFORMATION

Raikiri is a suspension concentrate containing the herbicide active ingredient mesotrione for the control of annual broad-leaved weeds in grain maize and forage maize crops only. Raikiri is absorbed by the weed foliage and works mainly by means of uptake through the leaves and limited uptake through the roots. After entry into the plant, mesotrione is translocated to the growing points. For best results, use when there is adequate soil moisture, under warm/humid conditions and when the weeds are small and actively growing. Following application sensitive weed species show chlorosis and subsequently die.

WARNINGS AND RESTRICTIONS

DO NOT USE on any crop that is being grown for seed.

DO NOT USE on any sweetcorn variety.

To avoid damage to plants outside the target area, DO NOT APPLY under windy conditions and take extreme care to avoid spray drift. All broad-leaved species are particularly susceptible to damage from spray drift. Do not apply to crops that are under stress from any cause including pest or disease attack, nutrient deficiency, drought, water-logging, previous herbicide application or any other cause.

Do not apply during very cold or frosty conditions or when large temperature fluctuations or excessive rainfall are forecasted or expected.

Do not allow spray bouts to overlap.

Before use, make sure the sprayer is cleaned from any previous use, and that it is calibrated to apply the recommended spray volume and pressure. Make sure that all the sprayer nozzles are performing equally and that the spray boom is set at the correct height above the crop.

Trace amounts of Raikiri remaining in the sprayer after use can cause damage to other crops subsequently treated with the same equipment. Immediately after use thoroughly clean the sprayer and all equipment as follows: fill the spray tank with clean water to a minimum of 10% of its total volume. Flush the water through the sprayer, including nozzles and spray booms and lines. Drain and repeat the procedure. During spray equipment cleaning, do not allow the product or sprayer equipment washings to contaminate water. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

CROPS AND TIMING

For use in grain maize and forage maize only. Apply from when the crop has between 2 to 8 leaves. For best results, use when there is adequate soil moisture, under warm/humid conditions and when the weeds are small and actively growing, see "WEEDS CONTROLLED"; for weed species and maximum sizes controlled. Ensure that the crop and weed foliage are dry at the time of application, do not use if

heavy rainfall is forecasted or expected soon after application. Under extreme weather conditions, some chlorosis (mild – moderate) may occur on the crop foliage. This chlorosis is normally transient and there is no effect on crop yield. Always check the condition of the crop and the weeds immediately prior to application. Treatment in poor growing conditions or in dry soil may give less reliable control.

RESISTANCE MANAGEMENT

Raikiri contains mesotrione which is from a group of herbicides referred to as 4-HPPD inhibitors. This class of compounds act by disrupting the synthesis of certain plant pigments involved in photosynthesis.

At the present time there is no known cross-resistance to mesotrione in weeds that exhibit resistance or reduced sensitivity to other herbicides with different modes of action. The use of Raikiri in mixtures and sequences with other herbicides approved for use in grain and forage maize crops and with a different mode of action is recommended to help to reduce the development of resistance. As herbicides in the group of 4-HPPD inhibitors are currently only available in maize crops, crop rotation with the use of herbicides with a different mode of action will also help to reduce the development of resistance. In the event that strains of weeds develop resistance or become less sensitive to a specific herbicide such as mesotrione, weed control may be reduced. Where maize crops are grown in successive seasons, avoid the use of Raikiri for more than two successive seasons to avoid the potential for weed resistance developing to mesotrione.

APPLICATION RATE

Apply Raikiri at 0.75-1.5 litres per ha in 200-300 litres of water per hectare. Select the dose rate from within this recommended range according to the weed species present, see "WEEDS CONTROLLED". Select the most appropriate water volume from within this recommended range to give good even cover of the weed foliage. Use the minimum water volume (200 litres per ha) wherever possible, but the higher volume (300 litres per ha) may be required if the weed foliage is dense. Do not exceed the maximum recommended water volume. Use a conventional hydraulic nozzle field crop sprayer with suitable nozzles operating at a pressure of 2-3 bar to apply a MEDIUM quality spray (BCPC). The use of any tank mix adjuvant with Raikiri is not required and is not recommended.

WEED CONTROL

Raikiri applied at 0.75 or 1.5 L/ha will control the broad leaf weed species listed in the table below, as well as Common barnyard grass, from emergence until the maximum growth stage as listed in the table:

WEED SPECIES	0.75 L/HA*	1.5 L/HA*	MAXIMUM GROWTH STAGE (BBCH) OR SIZE CONTROLLED
Black nightshade	S	S	10 leaves or 15cm
Charlock	S	S	Flowering
Common amaranth	S	S	8 leaves or 10cm
Common barnyard grass	-	MS	1 tiller
Common chickweed	S	S	10cm diameter
Fat hen	S	S	12 leaves or 20cm
Field pansy	S	S	6 leaves
Oilseed Rape (volunteer)	S	S	6 leaves
Prickly sow-thistle	MR	S	6 leaves
Redshank	MS	MS	6 leaves or 10cm

* S= Susceptible, MS = Moderately Susceptible, MR = Moderately Resistant

SAFETY PRECAUTIONS

OPERATOR PROTECTION

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVER ALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACE-SHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH SPLASHES from skin immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

Wash all protective clothing after use.

ENVIRONMENTAL PROTECTION

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms, respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

Raikiri

(MAPP 20553)

A suspension concentrate containing 100 g/litre mesotrione.
A selective herbicide for the control of annual broad-leaved weeds
and barnyard grass in grain and forage maize.

RISK & SAFETY INFORMATION

24 HOUR EMERGENCY NUMBER: +44(0)1763 212100

DANGER



- May cause an allergic skin reaction
 - Causes serious eye damage
 - Very toxic to aquatic life with long lasting effects
 - Suspected of damaging the unborn child
-
- Obtain special instructions before use.
 - Avoid breathing spray.
 - Wear protective gloves/protective clothing/eye protection/face protection.
 - IF ON SKIN: Wash with plenty of soap and water.
 - IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE or doctor/physician.
 - If skin irritation or rash occurs: Get medical advice/attention.
 - Collect spillage.
 - Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Raikiri – contains 100 g/litre (9.4% w/w) mesotrione.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT: The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work

Authorisation Holder

Sumi Agro Europe Ltd
68 Upper Thames St, London.
EC4V 3BJ

Marketing Company

SIPCAM UK Ltd, 4C Archway House,
The Lanterns, Melbourn Street,
Royston, Herts SG8 7BX

sipcamuk.com

Tel +44(0)1763 212100
Fax +44(0)1763 212101

Net contents:

5Le

Batch number: SEE PACKAGE



Scan to view the Safety Data Sheet

Alternatively, download the Safety Data Sheet
from sipcamuk.co.uk or contact your supplier.

PROTECT FROM FROST.
SHAKE WELL BEFORE USE.

UFI: EESN-9CM4-3A0V-YQA2

sipcamuk.co.uk

SIPCAM
UK

HERBICIDE

Raikiri

(MAPP 20553)

A suspension concentrate containing 100 g/litre mesotrione. A selective herbicide for the control of annual broad-leaved weeds and barnyard grass in grain and forage maize.

RISK & SAFETY INFORMATION
24 HOUR EMERGENCY NUMBER:
+44(0)1763 212100



Scan to view the Safety Data Sheet

Alternatively, download the Safety Data Sheet from sipcamuk.co.uk or contact your supplier.

Authorisation Holder

Sumi Agro Europe Ltd
68 Upper Thames St, London.
EC4V 3BJ

Marketing Company

SIPCAM UK Ltd, 4C Archway House,
The Lanterns, Melbourn Street,
Royston, Herts SG8 7BX

sipcamuk.co.uk

HERBICIDE

Net contents:
4x5Le

Batch number: SEE PACKAGE



Raikiri

(MAPP 20553)

A suspension concentrate containing 100 g/litre mesotrione. A selective herbicide for the control of annual broad-leaved weeds and barnyard grass in grain and forage maize.

RISK & SAFETY INFORMATION 24 HOUR EMERGENCY NUMBER: +44(0)1763 212100

GROUP 27 HERBICIDE

DANGER

- May cause an allergic skin reaction
- Causes serious eye damage
- Very toxic to aquatic life with long lasting effects
- Suspected of damaging the unborn child

- Obtain special instructions before use.
- Avoid breathing spray.
- Wear protective gloves/protective clothing/eye protection/face protection.
- IF ON SKIN: Wash with plenty of soap and water.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE or doctor/physician.
- If skin irritation or rash occurs: Get medical advice/attention.
- Collect spillage.
- Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Raikiri – contains 100 g/litre (9.4% w/w) mesotrione.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT: The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work

Authorisation Holder

Sumi Agro Europe Ltd, 68 Upper Thames St, London. EC4V 3BJ

Marketing Company

SIPCAM UK Ltd, 4C Archway House, The Lanterns, Melbourn Street, Royston, Herts SG8 7BX

sipcamuk.com

Tel +44(0)1763 212100

Fax +44(0)1763 212101



Scan to view the Safety Data Sheet

Alternatively, download the Safety Data Sheet from sipcamuk.co.uk or contact your supplier.

PROTECT FROM FROST.

SHAKEWELL BEFORE USE.

sipcamuk.co.uk



HERBICIDE

Net contents:

4x5Le

Batch number: SEE PACKAGE

