# Abyss

100% Seaweed Extract (Ascophyllum nodosum)

# Composition:

Raw material: Ascophyllum nodosum Organic carbon (C) 0.2% Mannitol 0.7g/L

Distributor:
Sipcam UK Ltd
4c Archway House,
The Lanterns
Melbourn Street,
Royston,
SG8 7BX

Tel. +44 (0) 1763 212100

Manufacturer: Sofbey SA Cso S.Gottardo 34 Chisso - 6830 CH Tel. +41 91 22 31 090

Abyss is a registered trademark of Sipcam Oxon S.P.A





**5**L

Always read the label

Keep out of reach of children.

Do not eat, drink or smoke when using this product.

Contains preservatives (biocides), complying with RD 506/2013 and regulation 528/2012.

If medical advice is needed, have the product container or label at hand. ABYSS is a Suspension Concentrate of 100% seaweed extract (*Ascophyllum nodosum*). Harvested from Irish coastal waters, ABYSS is processed within 24 hours using a vacuum, low temperature filtration to preserve the beneficial natural compounds and acidic pH. ABYSS contains a high concentration of natural compounds: carbohydrates (mannitol and alginates with chelating action), essential amino acids and vitamins to mitigate abiotic stress (stress, drought or heat) and enhance post-herbicide recovery. When used in fruit crops, the composition of ABYSS helps to promote flowering and fruit sets.

### DOSAGE AND RECOMMENDATIONS

CROPS	DOSE (L/HA)	RECOMMENDED GROWTH STAGE	NUMBER OF APPLICATIONS	INTERVAL
Cereals	0.5-1.0	From start of flowering	1-2	10-14 Days
Oilseed rape & Sunflower	0.5-1.0	From start of flowering	1-2	10-14 Days
Sugar beet	1.0	In combination with fungicide	3-4	10-15 Days
Outdoor leafy vegetables	1.0-1.5	In early development stages	2-3	7-10 Days
Outdoor horticultural crops	1.0-1.5	From pre-flowering to fruit growth	3-5	7-10 Days
Glasshouse horticultural crops	1.0-2.0	From pre-flowering to fruit growth	3-5	5-7 Days
Strawberries & Berries	1.0-1.5	From pre-flowering to fruits swelling	2-5	7-10 Days
Turf	1.0-2.0	Throughout vegetative cycle	-	7-14 Days
Ornamental plants	1.0-2.0	Throughout vegetative cycle	-	7-14 Days
Pome fruits	1.0-1.5	From pre-flowering to fruit growth	3-4	7-10 Days
Wine Grapes	1.0-2.0	From pre-flowering to pre-bunch closure stage	2-4	7-10 Days
Herb crops	1.0	Throughout vegetative cycle	-	7-14 Days
All crops during periods of stress	1.0-2.0	Preventative in case of frosts and to overcome stress from high/low temperatures. Damage from hail	2-4	2-7 Days

### MIXING ADVICE

ABYSS is compatible with crop protection products and/or fertilisers found on the market, but due to the wide variety of products, if mixing, first perform a compatibility jug test. When mixed with calcium-based products, first dissolve ABYSS in water (max concentration 20%) and then blend in the calcium-based product).

### STORAGE ADVICE

Store the product in a cool, dry and ventilated place. Protected from direct heat and sunlight. Protect against frosts. Shake well before use.

## OTHER INFORMATION

Caution: Only to be used according to the use and under the conditions set out in this label. Whoever uses the product is responsible for any damage resulting from an improper use.

Compliance with all the indications on this label is a condition to guarantee the effectiveness of the treatment and to avoid damage to plants, persons or animals. To avoid risks to humans or the environment, follow the instructions for use.

Dispose of the packaging in compliance with standards in force. The empty container must not be released into the environment and must not be reused.

### SIPCAM UK LTD CONDITIONS OF SUPPLY

All goods supplied by us are of high grade, but as we cannot exercise control over their mixing or use, all conditions or warranty, statutory or otherwise, as to the quality or fitness of our goods for any purpose are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.