

GAUNTLET®

CLETHODIM GUIDANCE



BEST PERFORMANCE



- Apply 0.7L/ha in a minimum of 200L/ha of water ensuring good coverage and uptake.
- Always apply with a water conditioner.
- Use drift reducing nozzles and apply in good spraying conditions.
- Use a suitable forward speed to ensure good coverage.
- Ensure grassweeds have reached 2-3 leaves and are actively growing.
- Require sufficient coverage of weed for optimum efficacy.
- Minimise spray overlap to reduce the risk of crop damage.
- Do not use ACCase inhibitors as the sole chemical method of control
- Use only as part of a resistance management strategy including cultural and chemical control.



TANK MIX



- Addition of water conditioner with a pH buffering system (no built in adjuvant) is recommended.
- Tank mixing with an insecticide for Cabbage Stem Flea Beetle control is permissible.
- Tank mixing with propyzamide is permissible for resistance management.
- Do not mix with propyzamide AND an insecticide.
- Do not mix with any adjuvants (Gauntlet has built in adjuvant)
- Do not mix with products other than those outlined above.



STEWARDSHIP



- Avoid applications after 15th October if any of the following criteria are met:
 - 6 leaf stage of the crop.
 - Early flowering or early maturing variety.
 - Crop drilled on or before 25th August.
- Do not apply after the end of October. No Spring applications.
- Avoid application of pesticide products 10 days prior or 14 days post use of clethodim.
- Nutritional products or insecticides may be used 7 days after clethodim application.



GUIDANCE



- Do not apply immediately after a period of frosty weather.
- Do not apply if foliage is wet.
- Do not apply in windy conditions.
- Ensure there is no crop stress or pest damage and crop waxing is adequate.

